arXiv:1304.5050v1 [math-ph] 18 Apr 2013

A note on the Jacobi identity

P.M. Lavrov [a], O.V. Radchenko [a] and I.V. Tyutin [b]

[a] *Tomsk State Pedagogical University,*
Kievskaya St. 60, 634061 Tomsk, Russia

[b] *I.E. Tamm Department of Theoretical Physics,*
P.N. Lebedev Physical Institute,
117924, Leninsky Prospect, 53, Moscow,Russia

Abstract

We present new basic identity for any associative algebra in terms of single commutator and anticommutators. From this identity we derive the set of four identities in terms of double commutators and anticommutators. Two of these four identities are independent. Equipping a given algebra with the basic identity we can prove that the multiplication in this algebra satisfies the associative condition. Extension of obtained results to supercase is found. A new identity for non-degenerate even symplectic (super)manifolds is proposed. Generalization of basic identity for the case of arbitrary number of elements is given.

Emails: lavrov@tspu.edu.ru, radchenko@tspu.edu.ru, tyutin@td.lpi.ac.ru

Keywords: associative algebras and superalgebras, Jacobi identity, symplectic supermanifolds

1 Introduction

Mathematical methods developed within Mathematical Analysis, Algebra, Differential Geometry, Differential Equations, Theory of Groups, Supermathematics and so on play a very important role in Modern Theoretical Physics. Formulations of Classical Mechanics and Classical Field Theory are based on symplectic manifolds endowed with the Poisson bracket satisfying the Jacobi identity [1, 2]. Quantum Mechanics intensively uses associative algebras of observable, differential equations, theory of groups, functional analysis [3, 4]. One of the main ingredients in consistent formulation of quantum mechanics is the commutator of operators which satisfies the Jacobi identity. Foundation of General Relativity is differential geometry on manifolds (see for instance [5]). The main object of General Relativity is the Riemann tensor which obeys in particular the Bianchi identity following from the Jacobi identity for covariant derivatives. Quantization of dynamical systems with constraints in Hamiltonian formalism [6, 7, 8, 9] assumes utilization of even symplectic supermanifolds [10] equipped with superextension of commutator and the Poisson bracket satisfying the generalized Jacobi identities. Quantum theory of gauge fields in the Lagrangian formalism is constructed with the help of odd symplectic supermanifolds endowed with the antibracket (the super Poisson bracket [11]) which satisfies the generalized Jacobi identity. List of such descriptions can be continued.

In present paper we are going to consider the Jacobi identity existing naturally for every associative algebra from a new point of view (in our knowledge). To do this we propose a new identity for any three elements of a given associative algebra using single commutators only instead of the Jacobi identity when double commutators are utilized. From this new identity the Jacobi identity can be derived but not on the contrary. Introducing the anticommutator we propose second identity for any associative algebra written for three elements of algebra in terms of single commutator and single anticommutators. We prove that the first identity written in terms of commutators follows from the second. It allows us speaking of this identity as a fundamental (basic) identity for any associative algebra. From the basic relation the four identities in terms of double commutators and anticommutators follow. It is proved that multiplication operation in an algebra equipped with the basic identity obeys the associative condition.

The paper is organized as follows. In Section 2, for any associative algebras we introduce new basic identity written in terms of single commutators and anticommutators and derive the set of four (reducible) identities in terms double commutators and anticommutators. We show that an algebra endowed with the basic identity is the associative algebra. In Section 3, we extend results obtained in previous section to supercase. In Section 4 we investigate a new identity existing for non-degenerate symplectic (super)manifolds. In Section 5, generalization of basic identities discussed in Section 2 is given. Finally, Section 6 gives concluding remarks.

2 New identities in associative algebras

Let us consider an associative algebra $\mathcal{A}$ with elements $X \in \mathcal{A}$. Let $T_i, i = 1, 2, ..., n$ be a basis in $\mathcal{A}$ so that for any element X of $\mathcal{A}$ there is the decomposition $X = x^i T_i$. Because $T_i T_j \in \mathcal{A}$ we have

$$T_i T_j = F_{ij}^{\ \ k} \, T_k \tag{2.1}$$

where $F_{ij}^{\ \ k}$ are structure constants. In terms of $F_{ij}^{\ \ k}$ the associative condition $(XY)Z = X(YZ)$ reads

$$F_{ij}^{\ \ n} F_{nk}^{\ \ \ m} = F_{j\,k}^{\ \ n} F_{in}^{\ \ \ m}. \tag{2.2}$$

We can always present these constants in the form

$$F_{ij}^{\ \ k} = \frac{1}{2} c_{ij}^{\ \ k} + \frac{1}{2} f_{ij}^{\ \ k} \tag{2.3}$$

where $c_{ij}^{\ \ k}$ and $f_{ij}^{\ \ k}$ obey the following symmetry

$$c_{ij}^{\ \ k} = c_{j\,i}^{\ \ k}, \quad f_{ij}^{\ \ k} = -f_{j\,i}^{\ \ k}. \tag{2.4}$$

Commutator $[\cdot, \cdot]$ and anticommutator $\{\cdot, \cdot\}$ are introduced for any two elements $X, Y \in \mathcal{A}$ by the relations

$$[X, Y] = XY - YX, \qquad \{X, Y\} = XY + YX, \tag{2.5}$$

which are elements of $\mathcal{A}$.

From (2.5) it follows the following Leibniz rules

$$[X, YZ] = [X, Y]Z + Y[X, Z], \qquad \{X, YZ\} = \{X, Y\}Z - Y[X, Z]. \tag{2.6}$$

It is clear that

$$[T_i, T_j] = f_{ij}^{\ \ k} T_k, \qquad \{T_i, T_j\} = c_{ij}^{\ \ k} T_k. \tag{2.7}$$

For any associative algebra $\mathcal{A}$ and for any $X, Y, Z \in \mathcal{A}$ without any reference to a basis the following identities exist

$$[X, YZ] + [Z, XY] + [Y, ZX] \equiv 0, \tag{2.8}$$

$$[X, YZ] + \{Y, ZX\} - \{Z, XY\} \equiv 0. \tag{2.9}$$

From these identities one can derive a set of identities in terms of double commutators and anticommutators. In particular, the Jacobi identity usually accepted in this case

$$[X, [Y, Z]] + [Z, [X, Y]] + [Y, [Z, X]] \equiv 0 \tag{2.10}$$

follows from (2.8). Moreover from (2.8) one can deduce the identity containing the anticommutator

$$[X,\{Y,Z\}]+[Z,\{X,Y\}]+[Y,\{Z,X\}]\equiv 0 \tag{2.11}$$

In similar manner from (2.9) one can derive the following identities

$$[X,\{Y,Z\}]-\{Z,[X,Y]\}+\{Y,[Z,X]\}\equiv 0, \tag{2.12}$$

$$[X,[Y,Z]]+\{Y,\{Z,X\}\}-\{Z,\{X,Y\}\}\equiv 0. \tag{2.13}$$

Note that the identities (2.8) and (2.9) are not independent because the summation of (2.9) over cyclic permutations gives the identity (2.8). Therefore we have the reason to consider the identity (2.9) as the fundamental identity in associative algebras because the identities (2.8), (2.10), (2.11), (2.12) and (2.13) can be derived from it. In its turn the set of identities (2.10)-(2.13) is not independent as well. Indeed, the summation of (2.12) over cyclic permutations gives the identity (2.11). The same operation applied to the identity (2.13) leads to the Jacobi identity (2.12) (see too [12]). Note that for associative algebras the identities (2.12), (2.13) were discussed in [13] as well. It is clear that having identities (2.12) and (2.13) one can reproduce the basic identity (2.9).

It should be noted that without any reference to a basis in $\mathcal{A}$ from the associative condition (2.2) one can derive analogs of identities (2.8)-(2.13) in terms of structure constants. In particular, the Jacobi identities for antisymmetric parts of structure constants, $f_{ij}{}^{k}$, have the form

$$f_{ij}{}^{m}f_{mk}{}^{n}+f_{ki}{}^{m}f_{mj}{}^{n}+f_{jk}{}^{m}f_{mi}{}^{n}\equiv 0. \tag{2.14}$$

The identities containing symmetric and antisymmetric parts of structure constants can be written as

$$c_{ij}{}^{m}f_{mk}{}^{n}+c_{ki}{}^{m}f_{mj}{}^{n}+c_{jk}{}^{m}f_{mi}{}^{n}\equiv 0, \tag{2.15}$$

$$f_{ij}{}^{m}c_{mk}{}^{n}-f_{ki}{}^{m}c_{mj}{}^{n}+c_{jk}{}^{m}f_{mi}{}^{n}\equiv 0, \tag{2.16}$$

$$c_{ij}{}^{m}c_{mk}{}^{n}-c_{ki}{}^{m}c_{mj}{}^{n}+f_{jk}{}^{m}f_{mi}{}^{n}\equiv 0. \tag{2.17}$$

Existence of the basic identity (2.9) allows us to discuss the associativity from a new point of view. Indeed, let $\mathcal{A}$ be an algebra. Then we can introduce the commutator and anticommutator by the rules (2.5) as elements of $\mathcal{A}$. Suppose the fulfilment of basic identity (2.9). We know that the identities (2.11)-(2.13) follow from (2.9). Then we can introduce the associative multiplication. To do this we define the multiplication " $\circ$ " for any two elements $X, Y \in \mathcal{A}$ by the rule

$$X\circ Y=\frac{1}{2}\Big([X,Y]+\{X,Y\}\Big). \tag{2.18}$$

In fact, we just put $X \circ Y = XY$. Let us check that the equality

$$(X \circ Y) \circ Z = X \circ (Y \circ Z) \tag{2.19}$$

holds. From (2.18) it follows

$$(X \circ Y) \circ Z = [[X,Y],Z] + [\{X,Y\},Z] + \{[X,Y],Z\} + \{\{X,Y\},Z\}, \tag{2.20}$$

$$X \circ (Y \circ Z) = [X,[Y,Z]] + [X,\{Y,Z\}] + \{X,[Y,Z]\} + \{X,\{Y,Z\}\}. \tag{2.21}$$

For the difference of (2.20) and (2.21) we obtain

$$\begin{aligned}(X \circ Y) \circ Z - X \circ (Y \circ Z) = -[Z,[X,Y]] - [X,[Y,Z]] - \\ -[Z,\{X,Y\}] - [X,\{Y,Z\}] + \{Z,[X,Y]\} - \{X,[Y,Z]\} + \\ +\{Z,\{X,Y\}\} - \{X,\{Y,Z\}\},\end{aligned} \tag{2.22}$$

or, taking into account the identity (2.13) and the Jacobi identity (2.10) which follows from it, the relation (2.22) reads

$$(X \circ Y) \circ Z - X \circ (Y \circ Z) = -[Z,\{X,Y\}] - [X,\{Y,Z\}] + \{Z,[X,Y]\} - \{X,[Y,Z]\}. \tag{2.23}$$

Now expressing $[Z,\{X,Y\}]$ and $[X,\{Y,Z\}]$ with the help of the identity (2.12) we arrive at the associativity (2.19). Therefore for any algebra $\mathcal{A}$ the basic identity (2.9) is equivalent to the associativity.

Any associative algebra obeys natural structure of the Lie algebra if one defines the Lie bracket through associative multiplication. The converse in general is not valid: the Lie bracket in general does not allow to introduce an associative multiplication. But if we consider an algebra $\mathcal{A}$ equipped with two bilinear operations $[\cdot,\cdot]$ and $\{\cdot,\cdot\}$ obeying the following symmetry properties:

$$[X,Y] = -[Y,X], \qquad \{X,Y\} = \{Y,X\}, \tag{2.24}$$

and satisfying the identities (2.12) and (2.13) then one can introduce the associative multiplication by the rule (2.18) (see as well [12]).

It should be stressed that consideration given above does not use assumptions for basis of algebra to be finite or infinite ones as well as to exist at all.

3 Superextension

Results obtained in the previous section can be extended to any associative superalgebras $\mathcal{A}$ with elements $X \in \mathcal{A}$ having the Grassmann parity $\varepsilon(X)$. Let $T_i, i = 1,2,...,n,\ \varepsilon(T_i) = \varepsilon_i$ be a basis in $\mathcal{A}$ so that for any element $X \in \mathcal{A}$ there is the decomposition $X = x^i T_i,\ \varepsilon(x^i) = \varepsilon(X) + \varepsilon_i$. Because $T_i T_j \in \mathcal{A}$ we have

$$T_i T_j = F_{ij}^{\ \ k}\, T_k \tag{3.1}$$

where $F_{ij}^{\ \ k}$, $(\varepsilon(F_{ij}^{\ \ k}) = \varepsilon_i + \varepsilon_j + \varepsilon_k)$ are structure constants. In terms of $F_{ij}^{\ \ k}$ the associativity $(XY)Z = X(YZ)$ reads

$$F_{ij}^{\ \ n} F_{nk}^{\ \ \ m} = F_{j\ k}^{\ n} F_{in}^{\ \ m} (-1)^{\varepsilon_i(\varepsilon_j + \varepsilon_k + \varepsilon_n)}. \tag{3.2}$$

In supercase the commutator $[\cdot,\cdot]$ and the anticommutator $\{\cdot,\cdot\}$ are introduced for any two elements $X, Y \in \mathcal{A}$ by the relations

$$[X, Y] = XY - (-1)^{\varepsilon(X)\varepsilon(Y)} YX, \qquad \{X, Y\} = XY + (-1)^{\varepsilon(X)\varepsilon(Y)} YX \tag{3.3}$$

with evident properties of symmetries

$$[X, Y] = -[Y, X](-1)^{\varepsilon(X)\varepsilon(Y)}, \qquad \{X, Y\} = \{Y, X\}(-1)^{\varepsilon(X)\varepsilon(Y)}. \tag{3.4}$$

From (3.3) the following Leibniz rules follow

$$[X, YZ] = [X, Y]Z + Y[X, Z](-1)^{\varepsilon(X)\varepsilon(Y)}, \tag{3.5}$$

$$\{X, YZ\} = \{X, Y\}Z - Y[X, Z](-1)^{\varepsilon(X)\varepsilon(Y)}. \tag{3.6}$$

For basis elements one has

$$[T_i, T_j] = f_{ij}^{\ \ k} T_k, \qquad \{T_i, T_j\} = c_{ij}^{\ \ k} T_k, \tag{3.7}$$

where structure constants $f_{ij}^{\ \ k}$ and $c_{ij}^{\ \ k}$ obey the following symmetry properties

$$c_{ij}^{\ \ k} = c_{ji}^{\ \ k}(-1)^{\varepsilon_i\varepsilon_j}, \quad f_{ij}^{\ \ k} = -f_{ji}^{\ \ k}(-1)^{\varepsilon_i\varepsilon_j}. \tag{3.8}$$

They can be identified with symmetric and antisymmetric parts of structure coefficients $F_{ij}^{\ \ k}$

$$c_{ij}^{\ \ k} = F_{ij}^{\ \ k} + F_{j\ i}^{\ k}(-1)^{\varepsilon_i\varepsilon_j}, \quad f_{ij}^{\ \ k} = F_{ij}^{\ \ k} - F_{j\ i}^{\ k}(-1)^{\varepsilon_i\varepsilon_j}. \tag{3.9}$$

For any associative superalgebra $\mathcal{A}$ and for any $X, Y, Z \in \mathcal{A}$ the following identities exist

$$[X, YZ](-1)^{\varepsilon(X)\varepsilon(Z)} + [Z, XY](-1)^{\varepsilon(Z)\varepsilon(Y)} + [Y, ZX](-1)^{\varepsilon(Y)\varepsilon(X)} \equiv 0 \tag{3.10}$$

$$[X, YZ](-1)^{\varepsilon(X)\varepsilon(Z)} + \{Y, ZX\}(-1)^{\varepsilon(Y)\varepsilon(X)} - \{Z, XY\}(-1)^{\varepsilon(Z)\varepsilon(Y)} \equiv 0. \tag{3.11}$$

These identities generalize relations (2.8) and (2.9). The identity (3.10) can be derived from (3.11) and (3.11) should again be considered as the basic identity for associative superalgebras. The set of identities in terms of double commutators and anticommutators reads

$$[X, [Y, Z]](-1)^{\varepsilon(X)\varepsilon(Z)} + [Z, [X, Y]](-1)^{\varepsilon(Z)\varepsilon(Y)} + [Y, [Z, X]](-1)^{\varepsilon(Y)\varepsilon(X)} \equiv 0, \tag{3.12}$$

$$[X,\{Y,Z\}](-1)^{\varepsilon(X)\varepsilon(Z)}+[Z,\{X,Y\}](-1)^{\varepsilon(Z)\varepsilon(Y)}+[Y,\{Z,X\}](-1)^{\varepsilon(Y)\varepsilon(X)}\equiv 0, \tag{3.13}$$

$$\begin{aligned}
&[X,\{Y,Z\}](-1)^{\varepsilon(X)\varepsilon(Z)}-\{Z,[X,Y]\}(-1)^{\varepsilon(Z)\varepsilon(Y)}+\\
&\qquad +\{Y,[Z,X]\}(-1)^{\varepsilon(Y)\varepsilon(X)}\equiv 0, && (3.14)\\
&[X,[Y,Z]](-1)^{\varepsilon(X)\varepsilon(Z)}+\{Y,\{Z,X\}\}(-1)^{\varepsilon(X)\varepsilon(Y)}-\\
&\qquad -\{Z,\{X,Y\}\}(-1)^{\varepsilon(Z)\varepsilon(Y)}\equiv 0. && (3.15)
\end{aligned}$$

The identities (3.12) and (3.13) follow from (3.15) and (3.14) respectively by the summations over cyclic superpermutations.

From the associativity (3.2) the identities in terms of symmetric and antisymmetric parts of structure constants follow

$$\begin{aligned}
&f_{ij}{}^{n}f_{nk}{}^{m}(-1)^{\varepsilon_i\varepsilon_k}+f_{ki}{}^{n}f_{nj}{}^{m}(-1)^{\varepsilon_k\varepsilon_j}+f_{j}{}^{n}{}_{k}f_{ni}{}^{m}(-1)^{\varepsilon_j\varepsilon_i}\equiv 0, && (3.16)\\
&c_{ij}{}^{n}f_{nk}{}^{m}(-1)^{\varepsilon_i\varepsilon_k}+c_{ki}{}^{n}f_{nj}{}^{m}(-1)^{\varepsilon_j\varepsilon_k}+c_{j}{}^{n}{}_{k}f_{ni}{}^{m}(-1)^{\varepsilon_i\varepsilon_j}\equiv 0, && (3.17)\\
&f_{ij}{}^{n}c_{nk}{}^{m}(-1)^{\varepsilon_i\varepsilon_k}-f_{ki}{}^{n}c_{nj}{}^{m}(-1)^{\varepsilon_k\varepsilon_j}+c_{j}{}^{n}{}_{k}f_{ni}{}^{m}(-1)^{\varepsilon_i\varepsilon_j}\equiv 0, && (3.18)\\
&c_{ij}{}^{n}c_{nk}{}^{m}(-1)^{\varepsilon_i\varepsilon_k}-c_{ki}{}^{n}c_{nj}{}^{m}(-1)^{\varepsilon_j\varepsilon_k}+f_{j}{}^{n}{}_{k}f_{ni}{}^{m}(-1)^{\varepsilon_i\varepsilon_j}\equiv 0. && (3.19)
\end{aligned}$$

We can again consider the associativity in superalgebras from a new point of view. Consider a superalgebra $\mathcal{A}$. Then we can introduce the commutator and anticommutator by the rule (3.3) as elements of $\mathcal{A}$. Suppose the fulfilment of basic identity (3.11). Introducing the multiplication for any two elements $X,Y\in\mathcal{A}$ by the rule

$$X\circ Y=\frac{1}{2}\Big([X,Y]+\{X,Y\}\Big) \tag{3.20}$$

and repeating the proof given in the previous section we arrive at the associativity of this multiplication

$$(X\circ Y)\circ Z=X\circ(Y\circ Z). \tag{3.21}$$

Any associative superalgebra obeys natural structure of the super Lie algebra if one defines the super Lie bracket through associative multiplication by formula $[X,Y]=XY-YX(-1)^{\varepsilon(X)\varepsilon(Y)}$. The converse in general is not valid. But if we can consider a superalgebra $\mathcal{A}$ equipped with two bilinear operations $[\cdot,\cdot]$ and $\{\cdot,\cdot\}$ obeying the following properties of generalized symmetry

$$[X,Y]=-(-1)^{\varepsilon(X)\varepsilon(Y)}[Y,X],\qquad \{X,Y\}=(-1)^{\varepsilon(X)\varepsilon(Y)}\{Y,X\}, \tag{3.22}$$

and satisfying the identities (3.14) and (3.15), then we can introduce an associative multiplication. Indeed, let us define the multiplication by the rule (3.20) and apply the proof given above. The we arrive at the associativity of this multiplication. In terms of this multiplication the binary operations introduced above have the usual representation

$$[X,Y]=X\circ Y-Y\circ X(-1)^{\varepsilon(X)\varepsilon(Y)},\quad \{X,Y\}=X\circ Y+Y\circ X(-1)^{\varepsilon(X)\varepsilon(Y)}. \tag{3.23}$$

Note that the proposal suggested above does not use any reference to a basis in a given superalgebra.

4 Non-degenerate symplectic (super)manifolds

Consider a non-degenerate symplectic supermanifold $(\mathcal{M},\omega)$ where $\mathcal{M}$ is a supermanifold and ω is a non-degenerate closed symplectic structure with Grassmann parity $\varepsilon(\omega(X,Y))=\varepsilon(X)+\varepsilon(Y)+\varepsilon(\omega)$ where X and Y are elements of the cotangent space of $\mathcal{M}$. We speak of even symplectic supermanifold when $\varepsilon(\omega)=0$ and of odd symplectic supermanifold if $\varepsilon(\omega)=1$. It is well-known (see for instance [1]) that an even symplectic supermanifold is the background underlying the description of classical dynamical systems in the Hamiltonian formalism. In its turn covariant quantization of gauge theories (Batalin-Vilkovisky method [11]) is based on a non-degenerate odd symplectic supermanifold. Any non-degenerate closed symplectic structure defines the super Poisson bracket $\{F,G\}$ $(\varepsilon(\{F,G\})=\varepsilon(F)+\varepsilon(G)+\varepsilon(\omega))$ which for any two scalar functions F, G on $\mathcal{M}$ is a scalar under general changes of coordinates on $\mathcal{M}$. The super Poisson bracket possesses the antisymmetry

$$\{F,G\}=-\{G,F\}(-1)^{(\varepsilon(F)+\varepsilon(\omega))\,(\varepsilon(G)+\varepsilon(\omega))}, \tag{4.1}$$

the linearity

$$\{F+G,H\}=\{F,H\}+\{G,H\}, \tag{4.2}$$

the Leibniz rule

$$\{F,GH\}=\{F,G\}H+\{F,H\}G(-1)^{\varepsilon(H)\varepsilon(G)}, \tag{4.3}$$

the Jacobi identity

$$\begin{aligned}&\{F,\{G,H\}\}(-1)^{(\varepsilon(F)+\varepsilon(\omega))\,(\varepsilon(H)+\varepsilon(\omega))}+\{H,\{F,G\}\}(-1)^{(\varepsilon(H)+\varepsilon(\omega))\,(\varepsilon(G)+\varepsilon(\omega))}+\\&+\{G,\{H,F\}\}(-1)^{(\varepsilon(G)+\varepsilon(\omega))\,(\varepsilon(F)+\varepsilon(\omega))}\equiv 0\end{aligned} \tag{4.4}$$

which is the consequence of closeness of symplectic structure.

In even case $(\varepsilon(\omega)=0)$ the super Poisson bracket coincide with superextension of the Poisson bracket. In odd case $(\varepsilon(\omega)=1)$ the super Poisson bracket presents the antibracket being one of the fundamental operations in the BV quantization method [11].

We want to note that in even case ($\varepsilon(\omega)=0$) from (4.1) and (4.3) it follows the identity

$$\{F,GH\}(-1)^{\varepsilon(F)\varepsilon(H)}+\{H,FG\}(-1)^{\varepsilon(H)\varepsilon(G)}+\{G,HF\}(-1)^{\varepsilon(G)\varepsilon(F)}\equiv 0. \tag{4.5}$$

Unfortunately, in contrast with associative algebras, we cannot consider this identity as a fundamental one for non-degenerate closed even symplectic supermanifolds because we cannot deduce the Jacobi identity (4.4) from (4.5). Nevertheless if the canonical quantization is applied to a dynamical system for which the phase space is described by a non-degenerate even closed symplectic supermanifold then the Poisson bracket is transformed into commutator $\{F,G\}\rightarrow (i\hbar)^{-1}[\hat{F},\hat{G}]$ and the identity (4.5) is reduced to (3.10) for operators $\hat{F},\hat{G},\hat{H}$.

For non-degenerate closed symplectic manifolds all Grassmann parities should be put into zero to obtain corresponding relations discussed above in this section.

5 Generalization of basic identities

Let us note that for any associative algebra the identities exist

$$[X_1,X_2X_3\cdots X_n]+cycle(X_1,X_2,...,X_n)\equiv 0,\qquad n=3,4,..., \tag{5.1}$$

which generalize the identity (2.8). The Jacobi identity

$$[[X_1,X_2],X_3]+[[X_3,X_1],X_2]+[[X_2,X_3],X_1]\equiv 0$$

follows from (5.1) for $n=3$

$$[X_1,X_2X_3]-[X_2,X_1X_3]+cycle(X_1,X_2,X_3)\equiv 0. \tag{5.2}$$

The generalized Jacobi identity for $n=4$ was discussed in [14] and had the form

$$[[[X_1,X_2],X_3],X_4]+[[[X_2,X_1],X_4],X_3]+[[[X_3,X_4],X_1],X_2]+[[[X_4,X_3],X_2],X_1]\equiv 0. \tag{5.3}$$

This identity can be derived from (5.1). Indeed, one can straightforwardly check that it is in fact

$$\begin{aligned}&[X_1,X_2X_3X_4]-[X_2,X_1X_3X_4]+[X_4,X_3X_2X_1]-[X_4,X_3X_1X_2]+\\&\quad +cycle(X_1,X_2,X_3,X_4)\equiv 0.\end{aligned} \tag{5.4}$$

The generalization of identity (5.3) for $n=5,6,...$ was given in [15] and again these generalized Jacobi identities can be derived from the fundamental identities (5.1).

We can also suggest a generalization of the identity (2.9). For four elements one has

$$[X_1,X_2X_3X_4]-\{X_4,X_1X_2X_3\}+\{X_3,X_4X_1X_2\}+[X_2,X_3X_4X_1]\equiv 0. \tag{5.5}$$

In general, for any associative algebra there exist the following identities

$$\begin{aligned}&[X_1, X_2X_3\cdots X_n] - \{X_n, X_1X_2\cdots X_{n-1}\} + \{X_{n-1}, X_nX_1\cdots X_{n-2}\} + \\ &+[X_{n-2}, X_{n-1}X_nX_1\cdots X_{n-3}] + [X_{n-3}, X_{n-2}X_{n-1}X_nX_1\cdots X_{n-4}] + \\ &+\cdots + [X_2, X_3\cdots X_nX_1] \equiv 0, \quad n \geq 4.\end{aligned} \tag{5.6}$$

The proof of (5.6) is based on the evident identity

$$\begin{aligned}&[X_n, X_1X_2\cdots X_{n-1}] + [X_{n-1}, X_nX_1\cdots X_{n-2}] + \\ &+\{X_n, X_1X_2\cdots X_{n-1}\} - \{X_{n-1}, X_nX_1\cdots X_{n-2}\} \equiv 0, \quad n \geq 2.\end{aligned} \tag{5.7}$$

Applying then to (5.7) the identity (5.1) we arrive at the identity (5.6). As in the case of (2.8) the identity (5.1) can be derive from (5.6). To do this we consider the summation of (5.6) over cyclic permutations. We have

$$\begin{aligned}&\big((n-2)[X_1, X_2X_3\cdots X_n] - \{X_n, X_1X_2\cdots X_{n-1}\} + \{X_{n-1}, X_nX_1\cdots X_{n-2}\}\big) + \\ &+cycle(X_1, X_2, ..., X_n) \equiv 0.\end{aligned} \tag{5.8}$$

But

$$\big(\{X_n, X_1X_2\cdots X_{n-1}\} - \{X_{n-1}, X_nX_1\cdots X_{n-2}\}\big) + cycle(X_1, X_2, ..., X_n) \equiv 0$$

and from (5.8) it follows the identity (5.1).

6 Discussions

We have introduced new basic identity for any associative algebra (2.9) and any associative superalgebra (3.11) and proposed to consider these identities as fundamental ones because they are described in terms of single commutators and anticommutators in contrast with usually discussed identities for algebras (see for instance [12, 13]).

We have proved that any algebra (superalgebra) equipped with the basic identity is in fact the associative algebra (superalgebra). It should be stressed that in the prove, there were no assumptions concerning finiteness or even existence of a basis for a given (super)algebra used.

We have introduced the identity (4.4) for any non-degenerate even symplectic supermanifold and discussed an application of this identity within canonical quantization of dynamical systems.

Finally we have proposed generalization of the basic identities for the case of arbitrary number of elements involving in these relations (see (5.1) and (5.6)).

Acknowledgments

The authors thank D. Leites and P. Zusmanovich for useful remarks and references concerning identities in algebras. The work on this paper was started during the visit of one (PML) of the authors to Juiz de Fora University, supported by the CAPES grant, project CAPES/PVE-1544128. The work of PML and OVR is supported by the LRSS grant 224.2012.2 and by the Ministry of Education and Science of Russian Federation, project 14.B37.21.0774. The work of PML is also supported by the RFBR grant 12-02-00121 and the RFBR-Ukraine grant 11-02-90445. The work of IVT is supported by the RFBR grant 11-01-00830.

References

[1] V.I. Arnold, *Mathematical Methods of Classical Mechanics* (Springer-Verlag, Berlin, 1978).

[2] N.N. Bogoliubov, D.V. Shirkov,*Introduction to the Theory of Quantized Field* (John Wiley and Sons Inc., 1959).

[3] P.A.M. Dirac, *The Principles of Quantum Mechanics* (Clarendon Press, Oxford, 1930).

[4] J. von Neumann, *Foundations of Quantum Mechanics* (Princeton University Press, 1955).

[5] R.M. Wald, *General Relativity* (The University of Chicago, 1984).

[6] I.A. Batalin I.A., Vilkovisky G.A., *Relativistic S-matrix of dynamical systems with boson and fermion constraints*, Phys. Lett. **69B** (1977) 309.

[7] I.A. Batalin I.A., Fradkin E.S., *A generalized canonical formalism and quantization of reducible gauge theories*, Phys. Lett., **122B** (1983) 157.

[8] D.M. Gitman, I.V. Tyutin, *Quantization of fields with constraints* (Springer, Berlin, 1990).

[9] M. Henneaux, C. Teitelboim, *Quantization of gauge systems* (Princeton U.P., Princeton, 1992).

[10] B. DeWitt, *Supermanifolds* (Second Edition, Cambridge University Press, 1992).

[11] I.A. Batalin, G.A. Vilkovisky, *Gauge algebra and quantization*, Phys. Lett. **B102** (1981) 27; *Quantization of gauge theories with linearly dependent generators*, Phys. Rev. **D28** (1983) 2567.

[12] M. Markl, E. Remm, *Algebras with one operation including Poisson and other Lie-admissible algebras*, J. Algebra **299** (2006) 171.

[13] J.-L. Loday, B. Vallette, *Algebraic Operados* (Springer, 2012).

[14] F. Wever, *Über Invarianten in Lie'schen Ringen*, Math. Ann. **120** (1949) 563.

[15] D. Blessnohl, H. Laue, *Generalized Jacobi identities*, Note di Matematika, Vol. **VIII** (1988) 111.